

**Comment on Phys.Rev.Lett.'s paper:
"Low temperature Electron Spin Resonance
of the Kondo Ion in a heavy fermion metal YbRh₂Si₂"**

R. J. Radwanski*

*Center of Solid State Physics, S^{nt} Filip 5, 31-150 Krakow, Poland,
@ Institute of Physics, Pedagogical University, 30-084 Krakow, Poland*

Z. Ropka

Center of Solid State Physics, S^{nt} Filip 5, 31-150 Krakow, Poland

The first successful Electron Spin Resonance (ESR) studies on single crystalline heavy fermion metallic compound YbRh₂Si₂, recently reported by Sichelschmidt *et al.* [1], has allowed for an unambiguous observation of the localized f states in this Kondo compound.

By this Comment [2] we would like to congratulate Prof. F. Steglich and his group for the first successful Electron Spin Resonance (ESR) on a heavy-fermion metallic compound at temperatures much below the Kondo temperature. ESR has been usually performed on a diluted ionic system - they have managed to observe it on the paramagnetic ion being the full part of a metallic solid. We fully agree with the interpretation of Ref. [1] that the revealed localized state is associated with the Kramers doublet of the Yb³⁺ configuration, to be more specific - with the $4f^{13}$ configuration of the Yb atoms.

Here we would like to report that the observed highly anisotropic g tensor with $g_{\perp} = 3.561$ and $g_{\parallel} = 0.17$ can be perfectly described by crystal-field (CEF) interactions allowing for a small off-tetragonal orthorhombic distortion. In the uniaxial CEF, whatever lower symmetry can be, there are only 4 Kramers doublets and in the tetragonal symmetry the ground state can be only a state Γ_6^1 or Γ_7^1 .

The perfect reproduction of the ESR results, $g_{\perp} = 3.561$ (with $g_L = 8/7$ it corresponds to $J_{\perp} = \pm 1.56$) and $g_{\parallel} = 0.17$ ($J_{\parallel} = \pm 0.08$) is obtained for the Γ_7^1 ground state for parameters: $B_2^0 = +14$ K, $B_4^0 = +60$ mK, $B_6^0 = -0.5$ mK, $B_4^4 = -2.30$ K and $B_6^4 = -10$ mK with a small local orthorhombic distortion $B_2^2 = +0.22$ K. These parameters yield the ground doublet:

$$\Gamma_7^1 = 0.803 \mid \pm 3/2 > + 0.595 \mid \mp 5/2 > - 0.026 \mid \mp 1/2 > - 0.008 \mid \pm 7/2 >$$

that is characterized by $J_{\perp} = \pm 1.560$, $J_{\parallel} = \pm 0.081$ and the quadrupolar operator value $Q_f =$

*URL: <http://www.css-physics.edu.pl>; Electronic address: sfradwan@cyf-kr.edu.pl

$3J_z^2 - J(J+1)$ of -4.7. The excited states are at 85 K (Γ_6^1 with $Q_f = -13.8$), 485 K (Γ_7^2) and 688 K (Γ_6^2). The admixture of last two small terms is an effect of the local orthorhombic distortion.

In case of the ground state Γ_6^1 the perfect reproduction of the experimental ESR results can be attained by:

$$\Gamma_6^1 = 0.944 | \pm 1/2 > + 0.322 | \mp 7/2 > - 0.052 | \mp 3/2 > - 0.046 | \pm 5/2 >$$

that yields $J_\perp = \pm 1.562$, $J_\parallel = \pm 0.083$ and Q_f of -11.2 This state can be obtained as the ground state by tetragonal CEF interactions $B_2^0 = +8$ K, $B_4^0 = -60$ mK, $B_6^0 = +6$ mK, $B_4^4 = -1.78$ K, $B_6^4 = -5$ mK with $B_2^2 = +0.475$ K, for instance. For these parameters the excited states are at 74 K (Γ_7^1 with $Q_f = -3.4$), 430 K (Γ_6^2) and 437 K (Γ_7^2). This state is less probable in comparison to the Γ_7^1 ground state as it gives too small anisotropy of the paramagnetic susceptibility $\chi(T)$.

The electronic structure with the Γ_7^1 ground state is very plausible, though we do not think that it is the final one. For its thermodynamical properties have to be more carefully analyzed, the best would be the direct inelastic-neutron-scattering experiment. However, apart of the g tensor the shown parameters reproduce the overall temperature dependence of the paramagnetic susceptibility $\chi(T)$ and its huge anisotropy, presented in Fig. 1a of Ref. [3], the preference for the magnetic ordering with moments perpendicular to the c axis, the magnetization curve for external magnetic fields up to 60 T applied along the tetragonal c axis (Fig. 1b of Ref. [3] - the magnetization at 2 K and at 60 T amounts to $0.85 \mu_B$). The derived electronic structure predicts a Schottky-type contribution to the specific heat with a maximum of 3.62 J/Kmol at about $T = 40$ K, superimposed on the lattice heat, and the anomalous temperature dependence of quadrupolar interactions. The excited doublet has much larger value of Q_f (in the absolute value) so with increasing temperature quadrupolar interactions, observed by means of the Mossbauer spectroscopy, should pass a maximum as it was discussed in Ref. [4]. Such the maximum is rather not expected in case of the Γ_6^1 ground state. A maximum in $Q_f(T)$ has been observed in some Yb compounds [5].

We do not think that the present set of CEF parameters is the final one. There is a plenty of sets that produce the shown ground-state eigenfunction (the simplest can be obtained by multiplication by a constant positive value) but surely the got set substantially confines the searching area of CEF parameters. We would like to express our big surprise that, despite of a persistent advocating by last 12 years for the localized picture for the heavy-fermion phenomena, so well-defined and so extremely thin atomic-scale energy levels exist in a metallic compound YbRh_2Si_2 though we have observed extremely thin energy levels in conventional rare-earth intermetallics [6, 7] and in a $3d$ Mott insulator [8].

In short, we welcome with the great pleasure the ESR results of Prof. F. Steglich group on heavy fermion metal YbRh_2Si_2 and we wish subsequent successful experiments on other Yb compounds. We fully agree with the F. Steglich's interpretation claiming the existence of the localized f electrons below Kondo temperature - this interpretation concurs with our long lasting claim that the evaluation of CEF interactions is indispensable for the physically adequate description of heavy-fermion compounds. We have derived CEF parameters of the tetragonal symmetry with a small orthorhombic distortion that perfectly reproduce the ESR values ($g_{\perp} = 3.561$ and $g_{\parallel} = 0.17$) as well as provide good reproduction of thermodynamical properties. The obtained parameters are surely not final as searching for a consistent set of CEF parameters is like a large puzzle but the discussion of heavy-fermion compounds within the same atomistic approach [7, 9] as conventional rare-earth compounds, where localized and band electrons coexist, is very plausible from the unification point of view. Also from a methodological point of view - the electronic structure and the importance of local distortions can be further experimentally verified. Finally, let express our feeling that the paper of Ref. [1] will be the turning point in the heavy-fermion subject.

-
- [1] J. Sichelschmidt, V.A. Ivanshin, J. Ferstl, C. Geibel and F. Steglich, Phys. Rev. Lett. **91**, 156401 (2003).
 - [2] due to the discriminating policy of the Editors of Phys.Rev.Lett. we can contribute to physics discussed in Phys.Rev.Lett. by means of Comments only.
 - [3] J. Custers, P. Gegenwart, C. Geibel, F. Steglich, T. Tayama, O. Trovarelli, and N. Harrison, Acta Phys. Pol. B **32**, 3211 (2001).
 - [4] R. J. Radwanski, J. Alloys-Compds **232**, L5 (1996).
 - [5] P. Bonville, J. Hammann, J.A. Hodges, P. Imbert, G. Jehanno, M.J. Besnus, and A. Meyer, Z. Phys. B: Condens. Matter **82**, 267 (1991).
 - [6] R. J. Radwanski, N.H. Kim-Ngan, F.E. Kayzel, J.J.M. Franse, D. Gignoux, D. Schmitt and F.Y. Zhang, J. Phys.: Condens. Matter **4**, 8853 (1992).
 - [7] R. J. Radwanski, R. Michalski, and Z. Ropka, Acta Phys. Pol. B **31**, 3079 (2000).
 - [8] Z. Ropka and R. J. Radwanski, Phys. Rev. B **67**, 172401 (2003).
 - [9] R. J. Radwanski and Z. Ropka, cond-mat/0010081.